



## Influence of microbial bioinoculants on growth and yield attributes of sugarcane

E Jamuna

Assistant Professor (Agricultural Microbiology), Oilseeds Research Station, Tamil Nadu Agricultural University, Tindivanam, Tamil Nadu, India

### Abstract

The study was conducted to evaluate the response of sugarcane variety CoC 24 to the application of bioinoculants viz., *Gluconoacetobacter diazotrophicus*, AM fungi and Azophos (*Azospirillum* and phosphobacteria), under different levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O inorganic fertilizer in plant and ratoon crop. The results revealed that the application of mycorrhizae, *G. diazotrophicus*, *Azospirillum* and phosphobacteria significantly produced higher cane yield in plant crop. The application of *Gluconoacetobacter diazotrophicus* @ 10 kg + AM fungi + Azophos @ 10 kg + 75 % NPK recorded the maximum germination and tiller population and also maximum mean millable cane population of 1.32 lakhs /ha, cane yield (137.45 t/ha) and sugar yield (16.96 t/ha). Similar results were recorded with the ratoon crop. The applied bacterial sources helps in nitrogen fixation and also in continuous mobilizing and solubilization of nutrients and their persistence and colonization in soil is an added advantage and also enhances the soil fertility. The usage of these bioinoculants in turn reduces the inorganic fertilizer input and thereby reduces the cost of cultivation.

**Keywords:** sugarcane: nitrogen fixing bacteria: phosphorous solubilizing: mobilizing: phosphobacteria: am fungi: yield parameters

### Introduction

Sugarcane is a very demanding crop as for a cane yield of 100 t/ha it removes about 205 kg N, 55 kg P<sub>2</sub>O<sub>5</sub>, 275 kg K<sub>2</sub>O and a large amount of micronutrients from soil (Yaduvanshi and Yadav, 1990) [16]. Since its fertilizer consumption is higher than that of other crops it has negative effect on soil health in the long term. In order to sustain productivity major nutrients are provided each year at the recommended application rates of 150 kg/ha of N and 60 kg each of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O for sugarcane. The efficiency of sugarcane to utilize N range between 16 and 45% as large quantities of applied N leach down through soil layer due to irrigation (Yadav and Prasad, 1992) [15]. Deterioration in the physico-chemical and biological properties of soil is considered to be the prime reason for declining sugarcane yield and productivity. The biofertilizer application increases crop growth through combination of BNF, growth promoting / hormonal substances, increased availability of soil nutrients and disease resistance. The importance of biofertilizer lies in the ability to supplement/ mobilize soil nutrients with minimal use of non-renewable resources. Johri (2006) [5] reported that some of the sugarcane varieties have been found to derive upto 70% of their nitrogen requirement through biological nitrogen fixation. Since then various kinds of bacteria such as *Glucanoacetobacter diazotrophicus*, *Herbaspirillum* spp., *Azospirillum amazonense*, *Burkholderia* spp., capable of fixing nitrogen have been reported to colonize the epidermis of sugarcane stem and roots of which *Glucanoacetobacter* seems to contribute substantially to nitrogen nutrition of the plant (Dobereiner *et al.*, 1995). *G. diazotrophicus* a nitrogen fixing endophyte is found in high number in all part of sugarcane (10<sup>2</sup> – 10<sup>6</sup> per g fresh weight) and its better colonization in sugarcane is probably due to this capability to grow in the presence of high sugar and low pH. Besides sugarcane it colonizes many other sugar and non-sugar crops

also. Production of plant growth hormones is the other beneficial trait associated with *G. diazotrophicus* (Sevilla *et al.*, 1998) [13]. Field trials conducted in sugarcane with *Glucanoacetobacter diazotrophicus* with other diazotrophs can match yield level equal to 275kg N/ha application (Sevilla *et al.*, 2001; Muthukumarasway *et al.*, 2002; Oliveria *et al.*, 2002) [12, 8, 10]. In India where sugarcane occupies over 4 million /ha, interest in exploitation of such microbes has increased due to an increase in prices of chemical fertilizers following withdrawal of subsidy on them. Concerning the above problems the current study was focused on the use of bioinoculants to enhance the sugarcane growth and also to assess the functional potentialities in relation to plant growth promoting activities like IAA, phosphate solubilisation and nitrogenase activity with the following objectives of standardizing the efficient combination of bioinoculants for maximizing sugarcane productivity and exploring the possibility of reduction in inorganic fertilizer input through bioinoculant application

### Materials and Methods

The experiment was conducted for a period of 2 year (2010-2012) at Sugarcane Research Station, Cuddalore with ten treatments with three replications in Randomised Block Design in plant and ratoon crop. The sugarcane variety taken for the study was CoC24. The mean maximum and minimum temperature of the location was 31.7 °C and 24.1 °C respectively. The mean annual rainfall was 1200 mm. The soil of the experimental field was sandy clay loam, with low available N (186.84 kg ha<sup>-1</sup>), medium in available 'P' (16.5 kg ha<sup>-1</sup>) and medium in available potash (265 kg ha<sup>-1</sup>). The pH of the soil is 7.2. The bioinoculants viz., *G. diazotrophicus*, AM fungi and Azophos (*Azospirillum* and phosphobacteria) was used along with inorganic fertilizer.

## Treatment details

**Table 1**

T1	:	<i>Gluconoacetobacter diazotrophicus</i> + 75 % NPK
T2	:	AM fungi (colonized root bits) + 75 % NPK
T3	:	Azophos + 75 % NPK
T4	:	<i>Gluconoacetobacter diazotrophicus</i> + AM fungi (colonized root bits) + 75 % NPK
T5	:	<i>Gluconoacetobacter diazotrophicus</i> + Azophos + 75 % NPK
T6	:	AM fungi (colonized root bits) + Azophos + 75 % NPK
T7	:	<i>Gluconoacetobacter diazotrophicus</i> + AM fungi (colonized root bits) + Azophos + 75 % NPK
T8	:	<i>Gluconoacetobacter diazotrophicus</i> + AM fungi (colonized root bits) + Azophos + 100% NPK
T9	:	Recommended NPK (100%) alone
T10	:	75% of recommended NPK alone

The microbial inoculants were applied as soil application @ 10 kg/ ha of *G. diazotrophicus* and Azophos, 25 kg of VAM / ha in two equal splits at 30<sup>th</sup> and 60<sup>th</sup> day after planting. The data collected on germination count, tiller population, millable cane population, cane yield commercial cane sugar per cent, sugar yield were pooled and analysed. The yield were recorded along with the quality parameters.

## Results and Discussion

The plant and the ratoon crop was raised during 2010-2012 with sugarcane variety CoC24. The microbial bioinoculants were applied as per the treatment schedule. The inorganic fertilizers were also applied as per schedule. Application of *G. diazotrophicus* @10 kg/ ha+ AM fungi @ 25 kg/ ha + Azophos @ 10 kg/ ha + 75% of the recommended NPK also recorded significantly maximum mean millable cane population and cane yield with 1.32 lakhs /ha and 137.45 t/ha respectively, followed by the application of *G. diazotrophicus* @10 kg + AM fungi @ 25 kg/ ha + Azophos @10 kg + 100% NPK with 1.29 lakhs millable cane / ha.

The CCS% and sugar yield also recorded maximum with 10.7% and 16.96 t/ha respectively with the application of *G. diazotrophicus* @10 kg/ ha+ AM fungi @ 25 kg/ ha + Azophos @ 10 kg/ ha + 75% of the recommended NPK (Table 1). As that of plant crop similar trend was observed with ratoon crop with maximum millable cane population of 1.22 lakhs/ha in plot treated with *G. diazotrophicus* @10 kg/ ha+ AM fungi @ 25 kg/ ha + Azophos @ 10 kg/ ha + 75% of the recommended NPK.

The increase in yield and enhanced quality parameters was due to the combined effect of the bioinoculants along with the inorganic fertilizers. The biofertilizers application enhanced the yield and quality parameters and also essential to maintain soil microflora population and protect soil fertility from deterioration. Significant changes in various plant growth parameters have been shown by the inoculation of various nitrogen fixing and plant growth promoting bacteria (Nayak *et al.*, 1986; Murty and Ladha, 1988 and Gunarto *et al.*, 1999)<sup>[9, 7, 3]</sup>. Sevilla *et al* (1998)<sup>[13]</sup>, have shown the benefits to sugarcane growth by using Nif – mutants of Acetobacter. In addition to nitrogen fixation the beneficial effects has been attributed to the production of plant growth hormones also (Glick, 1994)<sup>[2]</sup>. Application of phosphorous from different sources ie. From inorganic and as bioinoculants (AM fungi and phosphobacteria) was found to be effective in sugarcane. Continuous availability of the valuable nutrients and their persistence and colonization in soil makes the soil more fertile and healthy. The mobilization of P from soil to the plants is mediated by hairy root systems of the mycorrhizal fungi through plant roots. It commonly infect plant roots, including those of sugarcane forming beneficial symbiotic relationships (Kelly *et al.*, 1997)<sup>[6]</sup>. The improvement in plant growth was attributed to an enhanced access of mycorrhizal root to soil phosphorous located beyond the rhizosphere (Sanders and Tinker. 1973)<sup>[11]</sup> and infection by mycorrhizal fungi is significantly reduced at high soil phosphorous levels (Amijee *et al.*, 1989)<sup>[11]</sup>.

Mycorrhiza was found to be compatible with nitrogen fixers viz., Rhizobium (Hayman, 1986)<sup>[4]</sup>, Acetobacter and phosphate solubilising bacteria (Singh and Kapoor, 1999)<sup>[14]</sup>. It also holds good for sugarcane.

**Table 2:** Effect of combined application of bio inoculants with NPK fertilizers on yield attributes, juice quality, cane and sugar yield in plant crop

S.No.	Treatments	Millable cane population ('000/ha.)	Cane yield (' 000/ha.)	CCS%	Sugar yield (t/ha)
1.	<i>G. diazotrophicus</i> + 75% NPK	120.0	126.12	9.92	12.51
2.	AM fungi + 75% NPK	117.2	123.56	9.72	12.01
3.	Azophos + 75% NPK	118.7	125.79	9.80	12.33
4.	<i>G. diazotrophicus</i> + AM fungi + 75% NPK	124.8	131.25	10.34	13.57
5.	<i>G. diazotrophicus</i> + Azophos + 75% NPK	123.5	130.55	10.22	13.34
6.	AM fungi + Azophos + 75% NPK	121.2	129.84	9.95	12.92
7.	<i>G. diazotrophicus</i> + AM fungi + Azophos + 75% NPK	132.4	137.45	10.70	14.96
8.	<i>G. diazotrophicus</i> + AM fungi + Azophos + 100% NPK	125.2	133.62	10.50	14.03
9.	Recommended NPK (100%) alone	120.8	128.19	10.50	12.88
10.	75% of recommended NPK alone	95.70	107.15	10.05	10.52
	Mean	119.95	127.352	10.17	12.907
	SEd	4.21	3.55	0.99	0.68
	CD	8.47	7.14	NS	1.38

**Table 3:** Effect of combined application of bio inoculants with NPK fertilizers on yield attributes, juice quality, cane and sugar yield in ratoon crop

S. No.	Treatments	Millable cane population (' 000/ha.)	Cane yield (' 000/ha.)	CCS%	Sugar yield
1.	<i>G. diazotrophicus</i> + 75% NPK	116.20	118.42	10.37	12.05
2.	AM fungi + 75% NPK	114.25	115.86	10.17	11.55
3.	Azophos + 75% NPK	115.7	118.09	10.25	11.87
4.	<i>G. diazotrophicus</i> + AM fungi + 75% NPK	117.8	123.55	10.79	13.11

5.	<i>G. diazotrophicus</i> + Azophos + 75% NPK	117.0	122.85	10.67	12.88
6.	AM fungi + Azophos + 75% NPK	115.8	122.14	10.4	12.46
7.	<i>G. diazotrophicus</i> + AM fungi + Azophos + 75% NPK	122.4	129.75	11.15	14.5
8.	<i>G. diazotrophicus</i> + AM fungi + Azophos + 100% NPK	118.2	125.92	10.95	13.57
9.	Recommended NPK (100%) alone	116.5	120.49	10.95	12.42
10.	75% of recommended NPK alone	89.70	99.45	10.5	10.06
	Mean	114.36	119.65	10.62	12.45
	SEd	6.37	5.13	0.86	0.21
	CD	13.38	11.07	NS	0.58

## Conclusion

The inoculation of bioinoculants is beneficial for sugarcane growth for increasing the plant vigour at lower nitrogen levels, consequently the amount of fertilizer could be reduced. AM fungi and phosphobacteria are very much essential to convert the unavailable form of the phosphorous source to available source and providing to the plants. The usage of these bioinoculants inturn reduces the inorganic fertilizer input and thereby reduces the cost of cultivation. With this references these bioinoculants can be recommended for their use in nutrient management and enhanced sugarcane productivity.

## References

1. F Amijee, PB Tinker, DP Stribley. The development of endomycorrhizal root systems VII A detailed study of effects of soil phosphorous on colonisation. *New phytol.* 1989; 111:453-446.
2. BR Glick. The enhancement of plant growth by free living bacteria. *Canadian Journal of Microbiology.* 1994; 41:109-117.
3. Gunarto L, Adchi K, Senboku L. Isolation and selection of indigenous Azospirillum spp. from a subtropical island and effect if inoculation on growth of low land rice under several levels of N application. *Biol Fertile Soils.* 1999; 28:129-135.
4. Hayman H. Response of sugarcane to silicate source and rate growth and yield. *Phytopathology.* 1986; 93(4):402-412.
5. Johri H. Response of sugarcane to nitrogen source and rate of growth and yield. *Agron J.* 2006; 74:481-4.
6. Kelly RM, Edwards DG, Magarey RC, Thompson JP. The effect of VAM on the growth and nutrition of sugarcane. *Proc Aust Soc Sug cane Technol.* 1997; 19:73-79.
7. Murthy MG, Ladha LK. Influence of Azospirillum inoculation on the mineral uptake and growth of rice under hydroponic conditions. *Plant & Soil.* 1988; 108:281-285.
8. Muthukumarasamy R, Revathy G, Loganathan P. Effect of inorganic N on the population in vitro colonization and morphology of Acetobacter. *Plant & Soil.* 2002; 243:91-102.
9. Nayak DN, Ladha LK, Watanabe I. The fate of marker A. lipoferum inoculated and its effect on growth, yield and N fixation. *Biol Fertile Soils.* 1986; 2:7-14.
10. Oliveria ALM, Urquiaga S, Dobrenier J, Baldani JI. The effect of inoculating endophytic N fixing bacteria on micropropogated sugarcane plants. *Plant & Soil.* 2002; 242:205-215.
11. Sanders FE, Ticker PB. Phosphate flow into mycorrhizal roots. *Pesticide Science.* 1973; 4:385-395.
12. Sevilla M, Burris RH, Gunapala N, Kennedy C. Comparison of benefits of sugarcane plant growth with Acetobacter inoculation in wild type and Nif mutant strains. *Mol Plant Microbe Interaction.* 2001; 14:358-366.
13. Sevilla M, Oliveria D, Baldani I, Kenedy C. Contributions of the bacterial endophyte of Acetobacter to sugarcane nutrition: a preliminary study. *Symbiosis.* 1998; 25:181-191.
14. Singh, Kapoor. Silicon nutrition and sugarcane cultivation. *J Plant Nutr.* 1999; 15(7):1-10.
15. Yadav S, Prasad H. Nitrogenase activity of sugarcane propagated from stem cuttings on sterile vermiculite. *Soil Biol Biochem.* 1992; 12:413- 417.
16. Yaduvanshi, Yadhav. Isolation and characterisation of genetically two distant group of Acetobacter from new host plant. *J Appl Bacteriol.* 1990; 86:1053-1058.
17. Standford S, English L. Use of flame photometer in rapid soil tests of K and Ca. *Agron J.* 1949; 4:446-447.
18. Olsen SR, Cole F, Watanabe S, Dean AL. Estimation of available phosphorus in soil by extraction with sodium bicarbonate. *USDA. Circ.* 939, 1954.
19. Subbiah BV, Asija CL. A rapid procedure for estimation of available nitrogen in soils. *Curr Sci.* 1956; 25:259-260.
20. Omar EB Elnaeim, Yousif Ali Ragab. Esterification of cellulose from sugarcane bagasse by some long- chain acid chlorides. *International Journal of Chemistry Studies, Volume 4, Issue 2, 2020, Pages 28-32.*