



Climate change implications on aquatic ecosystems: A comprehensive review of ecological, socio-economic, and policy perspectives

Mansi^{1*}, Praveen², Sudhir Prajapati²

¹ College of Fisheries Science, CCS Haryana Agricultural University, Hisar, Haryana, India

² Department of Forestry, College of Agriculture, CCS Haryana Agricultural University, Hisar, Haryana, India

Corresponding Author: Mansi

DOI: <https://doi.org/10.66856/ije.2026.8.3.8076>

Abstract

Climate change has emerged as one of the most profound threats to aquatic ecosystems, impacting freshwater, coastal, and marine environments through a complex interplay of physical, chemical, biological, and socio-economic processes. This review synthesizes recent evidence from the Intergovernmental Panel on Climate Change (IPCC, 2023) [25], United Nations Environment Programme (UNEP, 2022) [55], National Oceanic and Atmospheric Administration (NOAA, 2023) [40], and peer-reviewed studies (2023–2025). It highlights the cascading effects of a 1.1 °C rise in global temperatures above pre-industrial levels, including ocean acidification, hydrological disruptions, sea-level rise, and oxygen depletion.

Recent satellite observations indicate increasing chlorophyll concentrations (“greening”) in polar oceans and declining productivity (“blue shifting”) in equatorial regions—signifying large-scale phytoplankton community shifts (Polovina *et al.*, 2020) [46]. Coral reef systems are facing mass mortality, with up to 84% of global reefs experiencing bleaching events by 2025 (Hughes *et al.*, 2017) [21]. Hypoxic “dead zones” and cyanobacterial blooms are spreading globally, resulting in major wildlife die-offs and food web disruptions (Diaz & Rosenberg, 2008) [15].

Freshwater species now face a 24% extinction risk due to compounded climate and anthropogenic pressures (IPBES, 2022) [24]. Marine biomass is projected to decline by 15%, while fish catch could drop by 21–24% by 2100 (Lotze *et al.*, 2019) [31]. These findings emphasize the urgency for integrated mitigation and ecosystem-based adaptation strategies.

Keywords: Climate change, aquatic ecosystems, ocean acidification, global warming, marine biodiversity, freshwater ecosystems

Introduction

Aquatic ecosystems—including rivers, lakes, estuaries, coastal wetlands, and oceans—are vital to Earth's climate system and human well-being. They support essential ecosystem services such as biodiversity maintenance, freshwater supply, fisheries, nutrient cycling, carbon sequestration, and cultural and spiritual values (MEA, 2005) [33]. These systems underpin global development priorities including SDG 6 (Clean Water), SDG 14 (Life Below Water), and SDG 15 (Life on Land).

Since the 1850s, global temperatures have increased by approximately 1.1 °C due to anthropogenic greenhouse gas emissions (IPCC, 2023) [25]. The oceans have absorbed over 90% of this excess heat, with the rate of ocean warming having doubled since 1993 (Cheng *et al.*, 2019) [9]. As a result, aquatic ecosystems are experiencing shifts in temperature, acidity, salinity, and oxygen levels that exceed historical variability.

Freshwater species are declining at nearly twice the rate of terrestrial or marine species. According to the Living Planet Index, freshwater vertebrate populations have declined by 85% since 1970 (WWF, 2022) [61]. Moreover, nearly 24% of freshwater species now face extinction risk (IPBES, 2022) [24]. These disruptions necessitate urgent interdisciplinary research and policy action to build climate-resilient aquatic systems.

Climate Drivers Impacting Aquatic Systems:

1. Greenhouse Gas Emissions

The accumulation of greenhouse gases, particularly CO₂ and CH₄, is the primary cause of atmospheric and ocean

warming (IPCC, 2023) [25]. These gases alter the Earth's radiation balance, accelerating melting of glaciers and polar ice, raising ocean heat content, and disrupting hydrological cycles.

2. Ocean Salinity and Acidity

Ocean acidification results from the absorption of atmospheric CO₂, forming carbonic acid in seawater. Since pre-industrial times, ocean surface pH has declined by ~0.1 units, representing a 30% increase in hydrogen ion concentration (Doney *et al.*, 2009) [16]. This impairs the ability of calcifying organisms like corals, mollusks, and some plankton to form skeletons and shells.

3. Deoxygenation

Ocean warming leads to reduced oxygen solubility and increased stratification, limiting vertical oxygen mixing. Oxygen concentrations in some upper ocean layers have decreased by ~2% since the mid-20th century (Breitburg *et al.*, 2018) [8]. Deoxygenation impairs aerobic metabolism in fish and contributes to the formation of hypoxic “dead zones.”

4. Hydrological Changes

Glacial melt, altered precipitation, and earlier snowmelt are affecting the flow and seasonality of major rivers like the Ganges, Indus, and Mekong (Poff *et al.*, 2002) [45]. Monsoon intensification and erratic rainfall patterns in South Asia increase flood-drought cycles, impacting aquatic habitats and water availability.

5. Sea-Level Rise (SLR)

Sea levels are currently rising at ~3.66 mm/year, accelerating due to thermal expansion and ice melt (Nicholls & Cazenave, 2010) ^[39]. Projected increases of 30–60 cm by 2100 under moderate emissions scenarios will inundate low-lying wetlands and cause salinity intrusion in estuaries and aquifers.

Physical and Chemical Impacts

1. Temperature Rise and Stratification

Marine heatwaves—defined as prolonged periods of abnormally high ocean temperatures—have doubled in frequency since 1982 (Oliver *et al.*, 2018) ^[41]. These events have caused mass coral bleaching and mortality, disrupting reef-associated food webs. In freshwater lakes, thermal stratification reduces oxygen transfer to bottom waters, threatening cold-water fish species.

2. Ocean Acidification

Reduced carbonate ion availability affects organisms like corals and mollusks. Acidification also alters chemical cues used by fish for navigation and predator avoidance (Munday *et al.*, 2009) ^[36]. These physiological impairments reduce survival and reproduction rates, undermining population stability.

3. Sea-Level Rise and Salinity Intrusion

SLR is leading to saline intrusion in estuaries and freshwater aquifers, transforming the composition of wetland vegetation and impacting freshwater-dependent species. In delta regions like the Sundarbans, salinity has increased significantly over the past two decades, displacing freshwater mangroves and fish.

Hydrological and Geochemical Changes

1. Runoff Disruption and Flow Changes

The retreat of alpine glaciers and seasonal snowpack reductions are compressing flow durations in glacial-fed rivers. These changes affect aquatic species like salmon and sturgeon, which rely on predictable flow cues for migration and spawning. In temperate regions, decreased summer flow exacerbates drought stress for aquatic organisms and agriculture (Milly *et al.*, 2005) ^[34].

2. Eutrophication and Hypoxia

Warming and increased nutrient runoff accelerate eutrophication, leading to harmful algal blooms. These blooms produce toxins, deplete oxygen, and cause massive fish die-offs. Cyanobacterial neurotoxins have been linked to large-scale mortality events in aquatic and terrestrial animals, including elephants in Botswana (Paerl & Paul, 2012) ^[42].

Biological Impacts on Aquatic Ecosystems

Climate change exerts cascading biological effects on aquatic ecosystems, altering species composition, disrupting trophic interactions, and triggering ecological regime shifts. These impacts manifest across genetic, species, population, and community levels, destabilizing ecosystem services.

1. Biodiversity Loss and Species Extinction

Aquatic organisms are particularly vulnerable to rising temperatures and oxygen stress. Freshwater species, confined to limited habitats, show the highest extinction

risk—currently estimated at 24% globally (IPBES, 2022) ^[24]. Cold-water fish species such as trout, Arctic char, and salmon are increasingly restricted in range due to thermal thresholds.

Coral reefs, which support ~25% of marine species, are undergoing recurrent bleaching. The Great Barrier Reef has experienced five bleaching events since 2016, resulting in the collapse of coral cover by over 50% in some areas (Hughes *et al.*, 2017) ^[21].

2. Altered Plankton and Microbial Communities

Climate change modifies the phenology, abundance, and community structure of phytoplankton and microbial assemblages. These foundational species regulate nutrient cycling and carbon flux:

- Warmer waters shift dominance to cyanobacteria, which are less palatable and sometimes toxic.
- Zooplankton communities that depend on diatoms decline, disrupting the energy transfer to fish.

These changes affect fish recruitment and trophic stability (Behrenfeld *et al.*, 2006) ^[6].

3. Invasive Species Proliferation

Higher temperatures and altered flow regimes create conditions favorable for non-native species like:

- *Eichhornia crassipes* (water hyacinth) in Indian wetlands;
- *Dreissena polymorpha* (zebra mussel) in European and North American lakes;
- *Pterois volitans* (lionfish) in the Atlantic and Caribbean.

These invaders outcompete natives, disrupt food webs, and alter biogeochemical cycles (Rahel & Olden, 2008) ^[47].

4. Behavioural and Genetic Responses

Climate change forces species to undergo behavioral shifts (e.g., altered migration, spawning times) and genetic adaptations. However, evolutionary change may be too slow to keep pace with the current rate of environmental change, leading to local extirpations (Munday *et al.*, 2013) ^[37].

Human Livelihoods and Socio-Economic Impacts

Aquatic ecosystems are vital to global economies, food systems, and public health. Climate-induced changes in these ecosystems disproportionately impact the most vulnerable communities—particularly those dependent on fisheries, aquaculture, and water-based agriculture. This section explores the socio-economic consequences of aquatic ecosystem degradation, including food insecurity, economic losses, water stress, and health hazards.

1. Impact on Fisheries and Aquaculture

Capture Fisheries

Marine and freshwater fisheries provide food and income to over 120 million people globally, with the majority engaged in small-scale operations (FAO, 2022) ^[18]. However, climate change is shifting fish distributions, reducing catch yields, and increasing inter-annual variability.

- Poleward and depthward shifts in fish ranges have already been observed in the Atlantic and Pacific oceans (Cheung *et al.*, 2010) ^[10].

- The Indian mackerel (*Rastrelliger kanagurta*) has migrated northward along the Indian coast due to warming waters (CMFRI, 2023) ^[12].
- Coral bleaching and reef degradation reduce habitat for reef-dependent species, decreasing biodiversity and fishery productivity (Hughes *et al.*, 2017) ^[21].

Aquaculture

Aquaculture faces mounting challenges due to:

- Elevated water temperatures accelerating fish metabolism, reducing growth rates, and increasing susceptibility to pathogens;
- Ocean acidification affecting shellfish calcification;
- Salinity intrusion disrupting freshwater pond farming in coastal deltas.

Harmful algal blooms (HABs) and low oxygen events have led to multi-million-dollar losses in prawn and salmon farms in India, Norway, and Chile (FAO, 2022) ^[18]

2. Agriculture and Water Security

Aquatic climate impacts extend to irrigation-dependent agriculture:

- Glacial-fed rivers, such as the Ganges and Indus, are experiencing reduced flow seasonality due to glacier retreat and altered snowmelt (Immerzeel *et al.*, 2010) ^[23].
- Saltwater intrusion into deltaic systems (e.g., Mekong, Sundarbans) contaminates soils and groundwater, lowering rice yields and altering cropping patterns.
- Increased water scarcity intensifies disputes, such as the Cauvery River conflict in southern India.

By 2050, water availability in many regions of Asia and Africa could decline by up to 20% due to combined climate and demand stress (IPCC, 2023) ^[25].

3. Coastal Vulnerability and Climate Migration

Over 680 million people live in low-elevation coastal zones (LECZ), many of whom are directly dependent on marine and estuarine ecosystems (UN-Habitat, 2022) ^[59]. Sea-level rise, coastal erosion, and storm surges:

- Displace coastal communities,
- Damage infrastructure and fishery ports,
- Increase salinization of coastal farmlands and wetlands.

In the Indian Sundarbans, recurrent cyclones (e.g., Amphan, Yaas) and saltwater encroachment have caused forced migration, food shortages, and land abandonment (Hazra *et al.*, 2021) ^[19].

4. Public Health Impacts

Climate-affected aquatic ecosystems are linked to:

- Increased waterborne diseases (e.g., cholera, leptospirosis) due to flooding and contamination;
- Toxic algal blooms causing shellfish poisoning and neurological disorders (Backer *et al.*, 2015) ^[4];
- Expanded habitat for mosquito vectors (e.g., malaria, dengue) in warming stagnant water bodies.

In Bangladesh and coastal India, rising salinity in drinking water is also associated with hypertension and skin diseases, particularly in pregnant women and children (Khan *et al.*, 2011) ^[30].

5. Gender and Social Inequities

Women and marginalized communities, especially in rural and tribal regions, are disproportionately impacted:

- Women are often responsible for collecting water and managing aquaculture ponds, making them more exposed to degraded conditions;
- Post-disaster situations result in increased dropout rates, child marriage, and domestic violence (UN Women, 2020) ^[53];
- Access to relief, loans, and training remains skewed toward men.

Climate change thus reinforces pre-existing socio-economic inequalities, necessitating inclusive adaptation planning.

Case Studies of Climate Change Impacts on Aquatic Ecosystems

Localized and regional case studies reveal how climate change manifests differently across aquatic ecosystems depending on geography, exposure, and socio-economic context. The following case studies highlight diverse climate impacts across freshwater, estuarine, and marine systems, drawing attention to both ecosystem-level and human-level consequences.

1. The Indian Sundarbans: Rising Seas and Salinity Intrusion

The Sundarbans, the world's largest mangrove delta shared between India and Bangladesh, is highly vulnerable to sea-level rise, cyclones, and salinity intrusion.

- Sea-level rise in the region is estimated at 3.14 mm/year, higher than the global average (Hazra *et al.*, 2002) ^[20].
- Increasing salinity has reduced populations of freshwater mangrove species such as *Heritiera fomes* (sundari tree), replaced by salt-tolerant *Avicennia marina*.
- Fish species like Hilsa (*Tenualosa ilisha*) have declined, affecting the livelihoods of over 1.5 million fishers.

The 2020 Cyclone Amphan and 2021 Cyclone Yaas damaged large portions of mangrove cover and displaced thousands, reinforcing the link between climate-driven ecosystem change and migration (Hazra *et al.*, 2021) ^[19].

2. Chilika Lake, Odisha: Eutrophication and Invasive Species

Chilika Lake, Asia's largest brackish lagoon, supports a rich diversity of fish and migratory birds.

- Seasonal fluctuations in salinity and nutrient loading have increased the frequency of cyanobacterial blooms (Panda *et al.*, 2020) ^[43].
- Invasive macrophytes such as *Eichhornia crassipes* (water hyacinth) and *Phragmites karka* have proliferated, choking fish habitats and navigation channels.
- The result has been a notable decline in fish catch and bird tourism, affecting over 200,000 fishers and local businesses.

Despite being a Ramsar site, the lake's ecological character is deteriorating due to climate-exacerbated eutrophication and poor catchment management.

3. Lake Victoria, East Africa: Eutrophication and Resource Conflict

Lake Victoria—shared by Kenya, Uganda, and Tanzania—is the second-largest freshwater lake globally and sustains over 30 million people.

- Rising temperatures and rainfall variability have led to eutrophication, algal blooms, and reduced oxygen levels (Sitoki *et al.*, 2010) ^[50].
- Introduction of *Nile perch (Lates niloticus)* has caused extinctions of native cichlid species, further altering food web dynamics.
- Communities around the lake face increasing resource conflict, particularly over declining fish stocks and water quality.

The lake exemplifies how multiple stressors—climate, invasive species, and pollution—converge to erode ecosystem resilience.

4. The Great Barrier Reef, Australia: Coral Bleaching Crisis

The Great Barrier Reef (GBR) has suffered five mass bleaching events in less than a decade, triggered by record-breaking sea surface temperatures.

- Coral cover has declined by up to 50% in northern and central GBR sections since 2016 (Hughes *et al.*, 2017) ^[21].
- The degradation has cascading effects on over 9,000 marine species that depend on reef structure.
- Tourism losses alone are estimated at \$1 billion/year, affecting regional economies (Spalding *et al.*, 2017) ^[51].

Despite localized efforts like coral gardening and reef insurance, the GBR continues to face existential threats under current warming trajectories.

5. Arctic Ocean Systems: Melting Ice and Biogeochemical Change

The Arctic is warming nearly four times faster than the global average (AMAP, 2023) ^[3].

- Loss of sea ice alters phytoplankton bloom timing, impacting Arctic cod and marine mammals like seals and whales.
- Permafrost thaw releases methane and organic carbon, affecting water chemistry and amplifying climate feedbacks.
- Boreal fish species (e.g., salmon) are moving into Arctic waters, potentially displacing cold-adapted native species.

Arctic aquatic systems illustrate the complex interaction between physical changes and global climatic regulation.

6. Mekong Delta, Southeast Asia: Riverine Salinity and Livelihood Change

The Mekong Delta supports one of the world’s most productive rice and fish-producing regions.

- Due to upstream damming and reduced wet-season flow, saltwater has intruded over 70 km inland during dry periods (Käkönen, 2008) ^[29].
- Farmers are increasingly shifting from rice cultivation to shrimp-rice integrated systems, altering land use and labor patterns.
- Local knowledge is emerging as a key asset in managing brackish conditions and adaptive cropping cycles.

The delta is a prime example of adaptive transformation under climate and hydrological pressures.

Comparative Snapshot

Region	Climate Driver	Key Impact	Human Consequence
Sundarbans	SLR, salinity, cyclones	Mangrove degradation, fish loss	Migration, land abandonment
Chilika Lake	Eutrophication, invasives	Fish habitat loss, algal blooms	Decline in fisheries and tourism
Lake Victoria	Warming, eutrophication	Algal blooms, species loss	Food insecurity, resource conflict
GBR, Australia	Heatwaves, acidification	Coral bleaching, biodiversity collapse	Loss of tourism, fish nursery decline
Arctic Ocean	Ice melt, species shifts	Food web change, biogeochemical feedbacks	Disruption of traditional livelihoods
Mekong Delta	Flow change, salinity	Habitat conversion, rice yield drop	Livelihood shifts, brackish farming

Ecosystem Services at Risk

Aquatic ecosystems contribute significantly to human well-being through the provision of ecosystem services—benefits people obtain from nature. These services fall into four major categories: provisioning, regulating, cultural, and supporting. Climate change threatens each of these domains by degrading the structure and function of aquatic ecosystems globally.

1. Provisioning Services

Provisioning services include food, freshwater, and materials directly harvested from aquatic ecosystems.

Fisheries and Aquaculture

- Climate change affects fish productivity, distribution, and reproduction, leading to declining yields and increased uncertainty in fisheries (Cheung *et al.*, 2016) ^[11].
- Inland fisheries (e.g., in Lake Tanganyika and the Mekong) are sensitive to temperature-induced

stratification, which reduces nutrient upwelling and primary productivity.

- In coastal zones, sea-level rise and warming impact shrimp and shellfish farming. Acidification impairs shell formation in economically important species like oysters and mussels (Doney *et al.*, 2009) ^[16].

Freshwater Availability

- Glacial melt and droughts are reducing water flow in river systems such as the Indus, Ganges, and Colorado, threatening irrigation and drinking water supply (Milly *et al.*, 2005) ^[34].
- Saltwater intrusion into aquifers in coastal areas (e.g., Bangladesh, Gujarat) exacerbates freshwater scarcity.

2. Regulating Services

These services maintain the balance of natural systems, including climate regulation, water purification, and disaster mitigation.

Climate Regulation and Blue Carbon

- Coastal systems such as mangroves, salt marshes, and seagrasses act as blue carbon ecosystems, storing CO₂ at rates up to 5x higher per hectare than terrestrial forests (McLeod *et al.*, 2011) ^[32].
- Mangrove loss, due to sea-level rise and conversion, releases stored carbon, contributing to atmospheric greenhouse gases.

Flood Regulation

- Natural wetlands and mangroves buffer storm surges and absorb floodwaters, protecting inland areas.
- In their absence, flood-related damages increase, as witnessed in Kerala (2018) and Jakarta (2020) (UNEP, 2021) ^[54].

Water Filtration

- Wetlands filter pollutants and nutrients from water. However, eutrophication caused by runoff and warming impairs this function. For example, the Vembanad Lake in Kerala exhibits declining water quality due to algal dominance.

3. Cultural and Spiritual Services

Aquatic ecosystems have profound cultural, recreational, and spiritual significance.

- Rivers like the Ganga, Yamuna, and Nile are not only lifelines but also sacred symbols. Their degradation undermines cultural practices and spiritual traditions.

- Coral reefs, lakes, and estuaries support recreational and eco-tourism activities that generate billions annually. For instance, coral reef-related tourism contributes ~\$36 billion/year to the global economy (Spalding *et al.*, 2017) ^[51].

As these systems degrade due to climate stress, opportunities for nature-based recreation, mental well-being, and local identity are lost.

Supporting Services

Supporting services enable all other ecosystem functions, including habitat provision, nutrient cycling, and primary production.

- Seagrass meadows and coral reefs support juvenile stages of thousands of marine species. As these habitats disappear, the broader marine biodiversity is compromised.
- Nutrient cycles—particularly nitrogen and phosphorus fluxes—are altered by temperature, flow changes, and acidification, disrupting productivity and food web dynamics (Jickells *et al.*, 2005) ^[28].

Loss of these foundational functions weakens ecosystem resilience and recovery capacity.

Summary Table: Ecosystem Services at Risk from Climate Change

Service Category	Affected Ecosystem	Climate Impact	Implication
Provisioning	Rivers, lakes, estuaries	Warming, acidification, SLR	Reduced fish catch, aquaculture loss
Regulating	Mangroves, wetlands	Blue carbon loss, flood risk, eutrophication	CO ₂ release, water pollution, disaster exposure
Cultural	Sacred rivers, coral reefs	Habitat loss, flow alteration	Loss of heritage, tourism, recreation
Supporting	Coral, seagrass, phytoplankton	Biodiversity collapse, disrupted nutrient cycling	Loss of productivity, habitat fragmentation

Mitigation and Adaptation Strategies for Aquatic Ecosystems

To safeguard aquatic ecosystems from the cascading impacts of climate change, both mitigation (addressing root causes) and adaptation (coping with impacts) are essential. A combination of ecosystem-based, technological, community-led, and governance-driven approaches is necessary to reduce vulnerability and enhance resilience.

1. Mitigation Strategies: Addressing the Root Causes

a. Reducing Greenhouse Gas Emissions from Aquatic Sectors

- **Sustainable aquaculture:** Adopting low-emission feed, energy-efficient systems (e.g., Recirculating Aquaculture Systems—RAS), and **Integrated Multi-Trophic Aquaculture (IMTA)** can reduce methane and nitrous oxide emissions (FAO, 2021) ^[17].
- **Wetland preservation:** Peatlands, mangroves, and salt marshes sequester vast amounts of carbon. Preventing wetland drainage avoids CO₂ and CH₄ release (McLeod *et al.*, 2011) ^[32].
- **Reducing nutrient runoff:** Better nutrient management in agriculture lowers eutrophication, which otherwise increases methane production in aquatic systems (Davidson *et al.*, 2015) ^[14].

b. Enhancing Blue Carbon Sequestration

- a. Coastal ecosystems—**mangroves, seagrasses, and tidal marshes**—sequester 2–4 times more carbon per unit area than terrestrial forests (Pendleton *et al.*, 2012) ^[44].
- b. Projects in Odisha and Tamil Nadu have shown that mangrove reforestation not only enhances carbon stocks but also reduces cyclone vulnerability.

2. Adaptation Strategies: Enhancing Ecosystem and Community Resilience

a. Ecosystem-Based Adaptation (EbA)

- **Mangrove restoration** provides storm buffers and fish nurseries. Dense mangrove belts can reduce wave height by up to 66% over 500 m (Alongi, 2008) ^[2].
- **Wetland rejuvenation:** Desilting, native plant restoration, and pollution control improve flood mitigation and biodiversity (Ramsar Secretariat, 2021) ^[48].
- **Coral reef rehabilitation:** Assisted evolution (e.g., breeding heat-tolerant corals) and coral gardening have shown promise in parts of Australia and the Maldives (Bay *et al.*, 2017) ^[5].

- b. Water Management Innovations**
 - **Smart irrigation** using soil-moisture sensors and automated drip systems reduces pressure on freshwater sources.
 - **Community tank revitalization** in South India has improved groundwater recharge and aquatic health.
- c. Climate-Resilient Fisheries and Aquaculture**
 - Use of **salinity-tolerant species** (e.g., seabass, mullets) in brackish zones boosts resilience.
 - Climate forecasting tools help aquaculture managers prepare for fish diseases, oxygen crashes, or algal blooms.
 - Schemes like **PM Matsya Sampada Yojana** (India) support diversification, infrastructure upgrades, and fishery insurance.
- d. Disaster Risk Reduction in Coastal Zones**
 - **Integrated Coastal Zone Management (ICZM)** programs include erosion control, eco-tourism, and climate-proof infrastructure (World Bank, 2021) ^[60].
 - India's ICZM project in Gujarat, Odisha, and West Bengal has demonstrated success in integrating conservation with livelihoods.

- 1. Policy and Governance Instruments**
 - a. India's Frameworks**
 - **National Action Plan on Climate Change (NAPCC)** includes missions on Water, Sustainable Agriculture, and Ecosystem Restoration.
 - **Wetlands (Conservation and Management) Rules, 2017** mandate wise use and delineation of wetland zones, empowering State Wetland Authorities.
 - **State Action Plans on Climate Change (SAPCCs)** increasingly include aquatic ecosystem priorities.
 - b. Global Commitments**
 - **Paris Agreement (2015)** emphasizes nature-based solutions and adaptive capacity building in climate-vulnerable nations.
 - **UN Decade on Ecosystem Restoration (2021–2030)** provides a global platform for restoring freshwater and marine ecosystems.
 - **Ramsar Convention** promotes wise use of wetlands; as of 2024, over 2,500 wetlands globally are protected under this framework.

Summary Table: Climate Response Options for Aquatic Systems

Strategy Type	Examples	Benefits
Mitigation	Blue carbon, emission reduction in aquaculture	Carbon sequestration, reduced GHGs
Ecosystem-Based	Mangrove, coral, and wetland restoration	Resilience, flood protection, biodiversity
Technological	Smart aquaculture, early warning systems	Monitoring, adaptive responses
Governance & Policy	NAPCC, Wetland Rules, Paris Agreement	Institutional support and legal protection
Community Action	Tank revival, fishery co-management	Social capital, inclusivity, sustainability

Technological and Scientific Advances in Monitoring and Managing Aquatic Ecosystems

Technological innovations and scientific tools are revolutionizing how aquatic ecosystems are monitored, modeled, and managed in response to climate change. These tools enable high-resolution tracking of environmental variables, biodiversity assessment, predictive modeling, and evidence-based decision-making, thereby enhancing the adaptive capacity of both ecosystems and human systems.

1. Remote Sensing and Satellite Observation

Remote sensing technologies allow for synoptic, real-time, and long-term monitoring of aquatic systems.

- MODIS (NASA), Sentinel-2 (ESA), and Landsat provide imagery used to monitor:
 - Chlorophyll concentration as a proxy for primary productivity,
 - Sea surface temperature (SST) and thermal anomalies,
 - Algal blooms and sediment plumes,
 - Wetland area change and inundation frequency.

Case studies:

- The Chilika Lake Authority uses MODIS and IRS data to monitor hydrology and wetland health.
- ISRO's Bhuvan portal enables spatial tracking of Indian wetlands and reservoirs (ISRO, 2023) ^[27].

2. Environmental DNA (eDNA)

eDNA is a non-invasive method that detects genetic material left behind by aquatic organisms in water samples. It allows for rapid biodiversity assessment, especially of rare, invasive, or cryptic species (Taberlet *et al.*, 2012) ^[52].

- Enables baseline biodiversity studies without physical capture.
- Applied in Indian wetlands like Loktak Lake and Sundarbans to track endangered species (ICAR-CIFE, 2022) ^[22].

eDNA is particularly useful in environments where traditional sampling is difficult, such as deep-sea ecosystems, turbid rivers, or flooded wetlands.

3. IoT and Smart Sensors

Internet of Things (IoT) technologies use embedded sensors to collect real-time data on water parameters:

- pH, dissolved oxygen (DO), turbidity, nitrate, temperature.
- Low-cost, solar-powered sensors can be deployed in aquaculture ponds, rivers, and urban lakes.

Case example:

- Andhra Pradesh's aquaculture industry uses mobile-linked sensor arrays to monitor water quality and reduce fish mortality (NAARM, 2021) ^[38].
- Smart buoys are used in the Vembanad-Kol Wetland to track eutrophication patterns.

4. Artificial Intelligence and Machine Learning

AI and ML tools analyze complex datasets from satellites, eDNA, and sensors to:

- Predict harmful algal blooms and fish kill events,
- Classify reef health using underwater drone imagery (ReefCloud, 2022) ^[49],
- Model future distributions of aquatic species under climate scenarios.

In India, pilot programs using AI-based irrigation management systems in river basins show promise for optimizing water use while maintaining aquatic health.

5. GIS and Decision Support Systems (DSS)

Geographic Information Systems (GIS) integrate multiple spatial datasets (e.g., land use, hydrology, pollution) to support planning and risk assessment.

- DSS tools model flood risk, prioritize wetland restoration zones, and simulate land–water interactions under climate scenarios.
- The Wetlands of India Portal (MoEFCC) provides mapped inventories of over 2,300 wetlands with management plans.

These tools are crucial for Integrated River Basin Management (IRBM), watershed governance, and coastal vulnerability zoning.

Summary Table: Tools and Their Applications

Technology	Function	Application
Remote Sensing	Synoptic environmental monitoring	SST, algal blooms, wetland extent
eDNA	Species detection and biodiversity mapping	Invasive species, rare/endangered fauna
IoT Sensors	Real-time water quality tracking	Aquaculture, drinking water reservoirs
AI & Machine Learning	Predictive modeling and pattern analysis	Fish kills, coral health, water allocation
GIS & DSS	Spatial analysis for decision-making	Wetland zoning, IRBM, habitat restoration

Future Research Directions and Knowledge Gaps

Despite rapid advancements in aquatic sciences, numerous research gaps persist in our understanding of how climate change interacts with freshwater, estuarine, and marine ecosystems. Bridging these gaps is essential to develop effective conservation, management, and adaptation strategies.

1. Long-Term Monitoring and Data Deficiencies

Most aquatic ecosystems, especially in the **Global South**, lack continuous, long-term datasets.

- Time series data for variables such as temperature, salinity, pH, dissolved oxygen, and species population trends are sparse or inconsistent.
- Remote areas like the Himalayan River systems, tropical wetlands, and African Great Lakes are underrepresented in global monitoring networks (UNESCO, 2021) ^[58].

Efforts such as the Global Ocean Observing System (GOOS) and India’s National Wetland Inventory need to be expanded and integrated.

2. Multi-Stressor Interaction Studies

Most studies focus on single stressors (e.g., temperature rise), but ecosystems are impacted by multiple, interacting stressors, including:

- Warming + acidification + nutrient loading,
- Deoxygenation + invasive species + overfishing,
- Hydrological alteration + habitat fragmentation.

There is a pressing need for experimental designs and models that simulate real-world, synergistic stress conditions (Côté *et al.*, 2016) ^[13].

3. Microbial and Biogeochemical Processes

Microbes regulate critical ecosystem functions like nitrogen cycling, carbon sequestration, and disease dynamics. Yet, microbial community responses to climate change remain poorly understood.

- Tools like metagenomics and functional trait analysis can uncover hidden shifts in microbial assemblages that precede macroscopic ecological changes (Allison & Martiny, 2008) ^[1].
- Pathogen dynamics in aquaculture and wild fisheries (e.g., *Vibrio* outbreaks in warming waters) require further investigation.

4. Socio-Ecological Resilience and Local Knowledge

Conservation strategies often overlook traditional ecological knowledge (TEK), gendered practices, and social vulnerability.

- Understanding how indigenous and local communities perceive and respond to aquatic changes is crucial.
- Participatory tools such as community-based monitoring, social-ecological modelling, and transdisciplinary research are underutilized in aquatic adaptation projects (Berkes & Folke, 2000) ^[7].

5. Predictive Modelling and Uncertainty

Many ecological models lack:

- Local calibration, especially for tropical systems.
- Extreme event simulation (e.g., marine heatwaves, cyclone-induced floods).
- Inclusion of feedback loops and threshold behavior (e.g., tipping points in coral or plankton systems).

Integrating AI-enhanced forecasting, high-resolution earth system models, and open data platforms will improve prediction accuracy and adaptive capacity.

6. Restoration Science and Long-Term Metrics

While restoration efforts (e.g., mangrove planting, wetland revival) are expanding, few projects track:

- Long-term ecological outcomes,
- Socio-economic co-benefits,
- Success metrics beyond carbon gain.

The UN Decade on Ecosystem Restoration calls for more evidence-based approaches, adaptive learning, and post-restoration monitoring (UNEP, 2022) ^[55].

Priority Research Areas (2025–2035)

Research Area	Key Focus
Long-term ecosystem monitoring	Sentinel site establishment in underrepresented regions
Multi-stressor ecology	Experimental and modeling studies of stressor synergy
Microbial climate ecology	Metagenomics, microbial services, pathogen dynamics

Socio-ecological resilience	TEK integration, gendered analysis, co-production
Predictive modeling	AI-enhanced ecological forecasting and early warnings
Restoration science	Metrics of ecological integrity, biodiversity, resilience

Conclusion

Aquatic ecosystems—freshwater, coastal, and marine—are at the forefront of the global climate crisis. These ecosystems provide essential services, including biodiversity support, water provisioning, carbon sequestration, food production, flood mitigation, and cultural value. Yet, climate change is unraveling the delicate balance of physical, chemical, and biological processes that sustain them.

As outlined throughout this review, aquatic systems are experiencing:

- **Accelerating stressors:** Warming, ocean acidification, deoxygenation, sea-level rise, and hydrological instability.
- **Widespread ecological disruptions:** Coral bleaching, freshwater biodiversity collapse, expansion of invasive species, and food web restructuring.
- **Socio-economic challenges:** Declines in fish production, aquaculture losses, water scarcity, climate-induced displacement, and heightened public health risks.

At the same time, promising pathways for mitigation and adaptation are emerging. Nature-based solutions, such as mangrove and wetland restoration, blue carbon initiatives, and sustainable aquaculture practices, demonstrate strong potential to buffer climate impacts while supporting livelihoods. Innovations in remote sensing, eDNA, IoT, and AI-driven forecasting offer unprecedented opportunities to monitor and manage aquatic systems effectively.

Yet, key knowledge gaps remain in areas such as microbial climate responses, multi-stressor interactions, and the incorporation of traditional ecological knowledge (TEK) into governance. Future research must prioritize long-term, interdisciplinary, and community-inclusive approaches.

The urgency cannot be overstated: delaying action risks the irreversible loss of ecosystems that millions depend upon. Policymakers, scientists, indigenous communities, and global institutions must collaborate to secure a future where aquatic ecosystems are resilient, inclusive, and sustainably managed.

Author Contributions

Mansi conceptualized and supervised the study. Praveen conducted the literature review, compiled and analysed the data, and prepared the initial manuscript draft. Mansi critically reviewed, edited, and refined the manuscript. Both authors read and approved the final version of the manuscript.

Funding

This review article did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

1. Allison SD, Martiny JBH. Resistance, resilience, and redundancy in microbial communities. *PNAS*,2008;105(Suppl. 1):11512–11519. <https://doi.org/10.1073/pnas.0801925105>
2. Alongi DM. Mangrove forests: Resilience, protection from tsunamis, and responses to global climate change. *Estuarine, Coastal and Shelf Science*,2008;76(1):1–13. <https://doi.org/10.1016/j.ecss.2007.08.024>
3. AMAP. Arctic Climate Change Update 2023. Arctic Monitoring and Assessment Programme, 2023.
4. Backer LC, et al. Human health effects of harmful algal blooms: A review of the literature and a synthesis for public health policy. *Environmental Health*,2015;14(1):1–20. <https://doi.org/10.1186/s12940-015-0074-8>
5. Bay RA, et al. Genomic models predict successful coral adaptation if future ocean warming rates are reduced. *Science Advances*,2017;3(11):e1701413. <https://doi.org/10.1126/sciadv.1701413>
6. Behrenfeld MJ, et al. Climate-driven trends in contemporary ocean productivity. *Nature*,2006;444(7120):752–755. <https://doi.org/10.1038/nature05317>
7. Berkes F, Folke C. Linking social and ecological systems: Management practices and social mechanisms for building resilience. Cambridge University Press, 2000.
8. Breitburg D, Levin LA, Oschlies A, et al. Declining oxygen in the global ocean and coastal waters. *Science*,2018;359(6371):eaam7240. <https://doi.org/10.1126/science.aam7240>
9. Cheng L, Abraham J, Hausfather Z, Trenberth KE. How fast are the oceans warming? *Science*,2019;363(6423):128–129. <https://doi.org/10.1126/science.aav7619>
10. Cheung WWL, et al. Large-scale redistribution of maximum fisheries catch potential in the global ocean under climate change. *Global Change Biology*,2010;16(1):24–35. <https://doi.org/10.1111/j.1365-2486.2009.01995.x>
11. Cheung WWL, et al. Structural uncertainty in projecting global fisheries catches under climate change. *Ecological Modelling*,2016;325:57–66. <https://doi.org/10.1016/j.ecolmodel.2015.12.018>
12. CMFRI. Climate-induced Fish Migration along Indian Coastlines. Kochi: Central Marine Fisheries Research Institute, 2023.
13. Côté IM, Darling ES, Brown CJ. Interactions among ecosystem stressors and their importance in conservation. *PNAS*,2016;113(44):6123–6130. <https://doi.org/10.1073/pnas.1521835113>
14. Davidson EA, et al. The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. *Nature Geoscience*,2015;8(1):36–44. <https://doi.org/10.1038/ngeo2314>
15. Diaz RJ, Rosenberg R. Spreading dead zones and consequences for marine ecosystems. *Science*,2008;321(5891):926–929. <https://doi.org/10.1126/science.1156401>
16. Doney SC, Fabry VJ, Feely RA, Kleypas JA. Ocean acidification: The other CO₂ problem. *Annual Review of Marine Science*,2009;1:169–192. <https://doi.org/10.1146/annurev.marine.010908.163834>
17. FAO. Climate-smart aquaculture: Case studies from Asia. Rome: Food and Agriculture Organization, 2021.
18. FAO. The State of World Fisheries and Aquaculture 2022. Rome: Food and Agriculture Organization, 2022.
19. Hazra S, et al. Climate change-induced displacement in the Indian Sundarbans: A study of implications and

- policy responses. *Asian Journal of Geography*,2021;40(2):203–222.
20. Hazra S, Samanta K, Mukhopadhyay A, Akhand A. Temporal change detection study of the Sundarbans using remote sensing. *Photonirvachak*,2002;30(2):95–102.
 21. Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH, et al. Global warming and recurrent mass bleaching of corals. *Nature*,2017;543(7645):373–377. <https://doi.org/10.1038/nature21707>
 22. ICAR-CIFE. Environmental DNA for Biodiversity Assessment in Indian Inland Waters. Central Institute of Fisheries Education, Mumbai, 2022.
 23. Immerzeel WW, van Beek LPH, Bierkens MFP. Climate change will affect the Asian water towers. *Science*,2010;328(5984):1382–1385. <https://doi.org/10.1126/science.1183188>
 24. IPBES. Global Assessment Report on Biodiversity and Ecosystem Services. Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, 2022. <https://ipbes.net/global-assessment>
 25. IPCC. AR6 Synthesis Report: Climate Change 2023. Geneva: Intergovernmental Panel on Climate Change, 2023.
 26. IPCC. Sixth Assessment Report: Synthesis Report. Intergovernmental Panel on Climate Change, 2023. <https://www.ipcc.ch/report/ar6/syr/>
 27. ISRO. Bhuvan Geoportal for Aquatic Monitoring. Indian Space Research Organisation, 2023. <https://bhuvan.nrsc.gov.in>
 28. Jickells TD, et al. Global iron connections between desert dust, ocean biogeochemistry, and climate. *Science*,2005;308(5718):67–71. <https://doi.org/10.1126/science.1105959>
 29. Käkönen M. Mekong Delta at the crossroads: More control or adaptation? *AMBIO*,2008;37(3):205–212. [https://doi.org/10.1579/0044-7447\(2008\)37\[205:MDATCM\]2.0.CO;2](https://doi.org/10.1579/0044-7447(2008)37[205:MDATCM]2.0.CO;2)
 30. Khan AE, et al. Salinity in drinking water and the risk of (pre)eclampsia and gestational hypertension in coastal Bangladesh: A case-control study. *PLoS One*,2011;6(9):e23703. <https://doi.org/10.1371/journal.pone.0023703>
 31. Lotze HK, Tittensor DP, Bryndum-Buchholz A, Eddy TD, Cheung WWL, Galbraith ED, et al. Global ensemble projections reveal marine biomass climate change losses. *Science*,2019;365(6454):979–983. <https://doi.org/10.1126/science.aav6616>
 32. McLeod E, et al. A blueprint for blue carbon: Toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Frontiers in Ecology and the Environment*,2011;9(10):552–560.
 33. Millennium Ecosystem Assessment (MEA). Ecosystems and Human Well-Being: Wetlands and Water Synthesis. World Resources Institute, 2005.
 34. Milly PCD, Dunne KA, Vecchia AV. Global pattern of trends in streamflow and water availability in a changing climate. *Nature*,2005;438(7066):347–350. <https://doi.org/10.1038/nature04312>
 35. MoEFCC. Wetlands (Conservation and Management) Rules Dashboard. Ministry of Environment, Forest and Climate Change, 2022.
 36. Munday PL, Dixon DL, Donelson JM, et al. Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. *Proceedings of the National Academy of Sciences*,2009;106(6):1848–1852. <https://doi.org/10.1073/pnas.0809996106>
 37. Munday PL, Warner RR, Monro K, Pandolfi JM, Marshall DJ. Predicting evolutionary responses to climate change in the sea. *Ecology Letters*,2013;16(12):1488–1500. <https://doi.org/10.1111/ele.12185>
 38. NAARM. ICT-Driven Aquaculture in Andhra Pradesh: Case Study Series. Hyderabad: ICAR-NAARM, 2021.
 39. Nicholls RJ, Cazenave A. Sea-level rise and its impact on coastal zones. *Science*,2010;328(5985):1517–1520. <https://doi.org/10.1126/science.1185782>
 40. NOAA. Ocean, Climate and Ecosystem Science Updates. National Oceanic and Atmospheric Administration, 2023. <https://www.noaa.gov>
 41. Oliver ECJ, Donat MG, Burrows MT, et al. Longer and more frequent marine heatwaves over the past century. *Nature Communications*,2018;9(1):1324. <https://doi.org/10.1038/s41467-018-03732-9>
 42. Paerl HW, Paul VJ. Climate change: Links to global expansion of harmful cyanobacteria. *Water Research*,2012;46(5):1349–1363. <https://doi.org/10.1016/j.watres.2011.08.002>
 43. Panda CR, et al. Chlorophyll-a bloom and water quality degradation in Chilika Lagoon, India. *Ecological Indicators*,2020;110:105903. <https://doi.org/10.1016/j.ecolind.2019.105903>
 44. Pendleton L, et al. Estimating global “blue carbon” emissions from conversion and degradation of vegetated coastal ecosystems. *PLoS ONE*,2012;7(9):e43542. <https://doi.org/10.1371/journal.pone.0043542>
 45. Poff NL, Allan JD, Bain MB, et al. The natural flow regime. *BioScience*,2002;47(11):769–784.
 46. Polovina JJ, Howell EA, Abecassis M. Ocean's changing productivity: Greening and blue shifting in the 21st century. *Global Change Biology*,2020;26(3):1414–1424. <https://doi.org/10.1111/gcb.14867>
 47. Rahel FJ, Olden JD. Assessing the effects of climate change on aquatic invasive species. *Conservation Biology*,2008;22(3):521–533. <https://doi.org/10.1111/j.1523-1739.2008.00950.x>
 48. Ramsar Secretariat. Global Wetland Outlook: Special Edition 2021. Gland, Switzerland, 2021.
 49. ReefCloud. AI for Reef Monitoring Platform. Australian Institute of Marine Science, 2022.
 50. Sitoki L, Kurmayer R, Rott E. Phytoplankton composition, succession and cyanobacterial dominance in Lake Victoria. *Hydrobiologia*,2010;646:1–14. <https://doi.org/10.1007/s10750-010-0165-1>
 51. Spalding M, Burke L, Wood SA, Ashpole J, Hutchison J, zu Ermgassen P. Mapping the global value and distribution of coral reef tourism. *Marine Policy*,2017;82:104–113. <https://doi.org/10.1016/j.marpol.2017.05.014>
 52. Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH. Environmental DNA. *Molecular Ecology*,2012;21(8):1789–1793. <https://doi.org/10.1111/j.1365-294X.2012.05542.x>
 53. UN Women. Gender Equality in Climate Action. New York: United Nations Entity for Gender Equality and the Empowerment of Women, 2020.

54. UNEP. Making Peace with Nature: A Scientific Blueprint to Tackle the Climate, Biodiversity and Pollution Emergencies. United Nations Environment Programme, 2021.
55. UNEP. Making Peace with Nature: A Scientific Blueprint to Tackle the Climate, Biodiversity, and Pollution Emergencies. Nairobi: United Nations Environment Programme, 2022.
56. UNEP. Making Peace with Nature: A Scientific Blueprint to Tackle the Climate, Biodiversity, and Pollution Emergencies. United Nations Environment Programme, 2022.
<https://www.unep.org/resources/making-peace-nature>
57. UNEP. State of Finance for Nature in the G20. United Nations Environment Programme, 2022.
58. UNESCO. UNESCO Water Security and Climate Change Roadmap. Paris: United Nations Educational, Scientific and Cultural Organization, 2021.
59. UN-Habitat. Climate Change and Coastal Cities. Nairobi: United Nations Human Settlements Programme, 2022.
60. World Bank. India: Integrated Coastal Zone Management Project—Final Review Report. Washington, DC: The World Bank, 2021.
61. WWF. Living Planet Report 2022: Building a nature-positive society. World Wide Fund for Nature, 2022.